

# AmproMed™ FOR CALVES

## (amprolium)

### 9.6% Oral Solution

### Coccidiostat



#### INDICATIONS

An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

#### DOSAGE AND ADMINISTRATION

5-Day Treatment Protocol	21-Day Prevention Protocol
Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are **96 mg** of Amprolium in every **1 mL** of AmproMed FOR CALVES

1 fl oz = 29.57 mL 1 fl oz = 2 measuring tablespoons	8 fl oz = ½ pint (236.56 mL) 16 fl oz = 1 pint (473.12 mL)
---	---

#### IN DRINKING WATER

Mix AmproMed FOR CALVES daily with fresh drinking water.

**5-DAY TREATMENT:** Add AmproMed FOR CALVES 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gallon. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AmproMed FOR CALVES 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

**AS A DRENCH:** AmproMed FOR CALVES drench solutions may be stored in a clean, closed labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz AmproMed FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of AmproMed FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

#### RESIDUE WARNINGS

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

#### WARNING

**NOT FOR HUMAN USE. Keep this and all drugs out of reach of children.**

#### PRECAUTIONS

**FOR ORAL USE IN ANIMLAS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

#### ADVERSE EVENTS

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse drug events, for technical assistance or to obtain a copy of the SDS contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

#### STORAGE

**Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F), with brief excursions to 40°C. Sodium Benzoate 0.1% add as preservative.**

#### Restricted Drug (California) – Use Only as Directed

Approved by FDA under ANADA # 200-482

# AmproMed™ FOR CALVES

(amprolium)  
9.6% Oral Solution  
Coccidiostat



List Number	Pack Size	Case Size
1AMP018	1 Gallon	4
1AMP027	16 fl oz	12

**MANUFACTURED BY:**

Bimeda, Inc.  
Le Sueur, MN 56058  
[www.bimeda.com](http://www.bimeda.com)

**N.A. CORP. ADDRESS:**

Bimeda, Inc.  
One Tower Lane, Oakbrook Terrace, IL 61081

For further information and queries, contact Bimeda at (888) 524-6332 or [US-info@Bimeda.com](mailto:US-info@Bimeda.com)

AmproMed™ is a trademark of Bimeda, Inc.  
Rev. 12/21 M-23-0355