

SpectoGard® Scour-Chek®

(spectinomycin)
50 mg per mL



ORAL SOLUTION FOR PIG SCOURS

Anti-Infective

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

EACH mL CONTAINS: Spectinomycin (from spectinomycin dihydrochloride pentahydrate) 50 mg. SpectoGard Scour-Chek contains spectinomycin. This antibiotic is fast-acting and effective against a variety of gram-negative and gram-positive organisms. SpectoGard Scour-Chek is a convenient dosage form of spectinomycin for baby pigs.

INDICATIONS

For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

WARNINGS

Residue Warnings

This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs.

Do not administer within 21 days of slaughter.

Human Warnings

NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN.

DOSE

Pigs under 10 lbs. – 1 pump (1 mL) twice daily.

Pigs over 10 lbs. – 2 pumps (2 mL) twice daily.

Each pump of the plunger delivers 1 mL of solution containing 50 mg of spectinomycin. Treatment may be continued twice daily for 3 to 5 days. If pigs do not improve within 48 hours, rediagnosis is suggested.

DIRECTIONS FOR USE

The plastic doser supplied in each 240 mL package with SpectoGard Scour-Chek makes treatment easy. After inserting the doser in the bottle, screw the top on tightly to avoid spilling. Push the piece of clear plastic tubing over the end of the pump. Then press the plunger a few times to fill the pump and the clear plastic tube with medication. To administer SpectoGard Scour-Chek, insert the plastic tube in the pig's mouth and press the plunger to obtain the desired dose.

When not in use remove the plunger from the bottle. Put the clear plastic tube in the neck of the bottle and press the plunger a few times to remove the medication from the pump. Replace the original cap on the bottle and always rinse out the doser with water to prevent sticking.

The 1000 mL bottle and 1 gallon bottle of SpectoGard Scour-Chek should be used only to refill the 240 mL bottle in the following manner.

1. Remove the cap and doser from the 240 mL bottle.
2. Unscrew the cap from the 1000 mL or 1 gallon bottle.
3. Pour, from the 1000 mL or 1 gallon bottle, the volume desired.
4. Reinsert the doser in the 240 mL bottle and screw cap on tightly.
5. Screw the cap tightly on the 1000 mL or 1 gallon bottle.

Follow the detailed instructions for use shown on the carton containing the 240 mL bottle of SpectoGard Scour-Chek.

HOW SUPPLIED

SpectoGard Scour-Chek is supplied in 240 mL multiple-dose plastic bottles. Each bottle is individually packed in a carton with a convenient plastic doser. The product is also supplied in 1000 mL resealable refills and 1 gallon plastic bottles to refill the 240 mL bottle.

STORAGE CONDITIONS

Store at controlled room temperature 20° - 25° C (68° - 77° F), with excursions permitted to 15° - 30° C (59° - 86° F).

SpectoGard® and Scour-Chek® are registered trademarks of Bimeda Inc.

Rev.09/22 M-23-0391

SpectoGard[®] Scour-Chek[®] (spectinomycin) 50 mg per mL



PRODUCT NO.	UNIT SIZE	CASE SIZE
1SPE022	240 mL	12
1SPE024	1000 mL	12
1SPE026	1 gallon	4

Approved by FDA under ANADA # 200-364

Restricted Drug (California) – Use Only as Directed

MANUFACTURED BY:

Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

N.A. CORP. ADDRESS:

Bimeda, Inc.
One Tower Lane
Oakbrook Terrace, IL 61081

For further information and queries, contact Bimeda at (888) 524-6332 or US-info@Bimeda.com

SpectoGard[®] and Scour-Check[®] are registered trademarks of Bimeda Inc.
Rev.09/22 M-23-0391