

SpectoGard[®]

(spectinomycin)

Water Soluble Powder

FOR USE IN GROWING CHICKENS

Anti-infective

NADA 038-661, Approved by FDA



MADE IN USA

INDICATIONS

SpectoGard[®] is indicated as an aid in the prevention or control of losses due to chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* (PPLO) and infectious synovitis associated with *Mycoplasma synoviae*.

Each 1000 g contains:

Spectinomycin dihydrochloride pentahydrate equivalent to spectinomycin activity 500.00 g

BENEFITS

- **Highly Soluble:** Mixes readily and thoroughly in water
- **Stable:** Effective over extended periods
- **Versatile:** Approved for chronic respiratory disease and infectious synovitis
- **Easy to Use:** Suitable for use in automatic water proportioners
- **Safe:** Approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1SPE027	1000 g (35.27 oz) jar	6

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

RESIDUE WARNINGS

DO NOT administer this drug within 5 days of slaughter.

DO NOT administer to laying chickens.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: Sales@BimedaUS.com

See reverse side for additional information and Dosage



SpectoGard®

(spectinomycin)

Water Soluble Powder

FOR USE IN GROWING CHICKENS

Anti-Infective

NADA 038-661, Approved by FDA

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) – Use only as directed.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

SpectoGard® is a water soluble powder intended as an anti-infective for growing chickens.

Each 1000 gram bottle contains spectinomycin dihydrochloride pentahydrate equivalent to 500 grams of spectinomycin activity.

ACTIVE INGREDIENTS:

Each gram contains spectinomycin dihydrochloride pentahydrate equivalent to 0.5 grams of spectinomycin activity.

INDICATIONS, DOSAGE & ADMINISTRATION:

As an aid in the prevention or control of losses due to CRD associated with *M. gallisepticum* (PPL0). Administer two (2) grams per gallon of water as the only source of drinking water for the first three days of life and for one day following each vaccination.

As an aid in controlling infectious synovitis due to *Mycoplasma synoviae* in broiler chickens. Administer one (1) gram per gallon of water as the only source of drinking water for the first three to five days of life.

DIRECTIONS FOR USE:

To provide 2 g/gal: Dissolve contents in 250 gallons of water. For proportioners, dissolve contents of 1/2 package in 1 gallon of water. Meter at 1 oz/gal.

To provide 1 g/gal: Dissolve contents in 500 gallons of water. For proportioners, dissolve contents of 1/2 package in 2 gallons of water. Meter at 1 oz/gal.

PRECAUTION:

Rarely, some individuals who handle spectinomycin develop serious reactions involving skin, nails and eyes. Individuals who have experienced rash or other evidence of allergic reaction should avoid further contact with spectinomycin.

RESIDUE WARNING:

DO NOT administer this drug within 5 days of slaughter.
DO NOT administer to laying chickens.

HOW SUPPLIED:

SpectoGard® is supplied in 1000 g (35.27 oz) screw cap, plastic bottles packed 6 per case.

STORAGE:

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

To obtain an SDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.