

# AmproMed™ FOR CALVES

**(amprolium)**


**20% Soluble Powder/Coccidiostat**

## INDICATIONS

An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

## BENEFITS

- **Effective:** breaks up the coccidia life cycle
- **Versatile:** labeled specifically for both the prevention and treatment of coccidiosis in beef and dairy calves
- **Short withdrawal:** only 24 hours after the last administration
- **Convenient:** treat the whole herd through the drinking water
- **Economical:** a single 10 oz pack treats 25 calves weighing 100 lb on the 5-day treatment regimen
- **Palatable:** accepted by calves in the drinking water
- **Safe:** approved by FDA



## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1AMP006	10 oz (283.5 g) pouch	2 x 12 x 10 oz pouches

See reverse side for Administration and Dosage.

**Bimeda, Inc.**  
One Tower Lane, Suite 2250  
Oakbrook Terrace, IL 60181 USA  
Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035  
Email: [sales@bimedaus.com](mailto:sales@bimedaus.com)

[www.bimedaus.com](http://www.bimedaus.com)



# AmproMed™ FOR CALVES

(amprolium)

20% Soluble Powder/Coccidiostat

ANADA 200-464, Approved by FDA

**FOR ANIMAL USE ONLY**

**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

**Restricted Drug (California) – Use Only as Directed**

**INDICATIONS:**

An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

**USE DIRECTIONS:**

There are **200 mg** of amprolium in every **1 gram** of AMPROMED FOR CALVES.

**5-Day Treatment Protocol**

Daily Dosage: 10 mg amprolium/kg  
(10 mg per 2.2 lb body weight)  
1 oz = 3 ½ measuring tablespoons

**21-Day Prevention Protocol**

Daily Dosage: 5 mg amprolium/kg  
(5 mg per 2.2 lb body weight)  
1 oz = 28.35 grams

**In Drinking Water:** Mix AMPROMED FOR CALVES daily with fresh drinking water. Stock solution for water proportioners may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:**

Add AMPROMED FOR CALVES to drinking water at the rate of 4 oz/50 gallon. One 10 oz pack to 125 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1 pouch (10 oz) in 1 gallon of water to make stock solution. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:**

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AMPROMED FOR CALVES to drinking water at the rate of 4 oz/100 gallon. One 10 oz pack to 250 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add ½ pouch (5 oz) in 1 gallon of water to make stock solution. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

**As a Drench:** AMPROMED FOR CALVES drench solutions may be stored in a clean, labeled container for up to 3 days.

**5-DAY TREATMENT:**

Add 3 oz AMPROMED FOR CALVES to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Treatment should be on a herd basis only: when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:**

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 ½ oz AMPROMED FOR CALVES to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**CAUTION:**

For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**RESIDUE WARNING:**

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS contact Bimeda, Inc. at 1-888-524-6332.

**STORAGE:**

Store at 20°C-25°C (68°F-77°F), excursions permitted between 15°C-30°C (59°F-86°F).

**HOW SUPPLIED:** 10 oz (283.5 g) Pouches.

