

AmproMed™ FOR CALVES

(amprolium)

9.6% Oral Solution/Coccidiostat

ANADA 200-482, Approved by FDA



MADE IN USA

INDICATIONS

An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves.

BENEFITS

- **Effective:** disrupts the coccidia life cycle
- **Versatile:** approved for both the prevention and treatment of coccidiosis in beef and dairy calves
- **Short withdrawal:** only 24 hours after the last administration
- **Economical:** one gallon of concentrate treats 1,600 gallons of drinking water based on a 21-day prevention regimen
- **Convenient:** treat the entire herd through the drinking water
- **Easy to mix:** concentrated solution disperses readily into the drinking water
- **Safe:** approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1AMP018	1 gallon (128 fl oz) (3.785 L)	4 x 1 gallon (512 fl oz) (15.141 L)
1AMP027	16 fl oz (473 mL)	12 x 16 fl oz

RESIDUE WARNING

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

See reverse side for Administration and Dosage.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com



AmproMed™ FOR CALVES

(amprolium)

9.6% Oral Solution/Coccidiostat

ANADA 200-482, Approved by FDA

FOR ORAL USE IN ANIMALS ONLY

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN

Restricted Drug (California) – Use Only as Directed

INDICATIONS:

An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves.

For a satisfactory diagnosis, a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

USE DIRECTIONS:

There are **96 mg** of amprolium in every **1 mL** of AmproMed™ FOR CALVES.

5-Day Treatment Protocol

Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

1 fl oz = 29.57 mL

1 fl oz = 2 measuring tablespoons

21-Day Prevention Protocol

Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

8 fl oz = ½ pint (236.56 mL)

16 fl oz = 1 pint (473.12 mL)

IN DRINKING WATER:

Mix AmproMed™ FOR CALVES daily with fresh drinking water.

5-DAY TREATMENT:

Add AmproMed™ FOR CALVES to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION:

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AmproMed™ FOR CALVES to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH:

AmproMed™ FOR CALVES drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT:

Add 3 fl oz AmproMed™ FOR CALVES to 1 pt (16 oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION:

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1½ fl oz of AmproMed™ FOR CALVES to 1 pt (16 fl oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

RESIDUE WARNING:

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING:

Not for Human Use. Keep this and all drugs out of reach of children.

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS, contact Bimeda at 1-888-524-6332.

PRECAUTIONS:

MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention. **Sodium Benzoate 0.1% added as preservative.**

STORAGE:

Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F), with brief excursions to 40°C.

