

Flunazine®-S

(flunixin meglumine)

Injectable Solution

50 mg/mL • Veterinary
FOR USE IN SWINE

ANADA 200-489, Approved by FDA

INDICATIONS

Flunazine®-S (*flunixin meglumine*) Injectable Solution is indicated for the control of pyrexia associated with swine respiratory disease.

BENEFITS

- **Swine claim:** a flunixin meglumine injectable approved for use in swine
- **Reduces fever:** pigs feel better which results in better performance
- **Fast acting:** maximum plasma concentrations in as little as 0.4 hours post administration
- **Long lasting:** quantifiable drug concentration can be measured up to 18 hours following IM injection
- **Potent:** more relief than with pentazocine, meperidine and codeine
- **Safe:** Non-narcotic, non-steroidal; no known contraindications when used as directed



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1FLU009	100 mL	12

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

RESIDUE WARNINGS

Swine must not be slaughtered for human consumption within 12 days of the last treatment.

Bimeda, Inc.

One Tower Lane, Suite 2250
Oakbrook Terrace, IL 60181 USA
Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035
Email: sales@bimedaus.com

See reverse side for Administration and Dosage.



Flunazine[®]-S

(flunixin meglumine)

Injectable Solution

50 mg/mL • Veterinary
FOR USE IN SWINE

ANADA 200-489, Approved by FDA

FOR ANIMAL USE ONLY
NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

FOR INTRAMUSCULAR USE IN SWINE
NOT FOR USE IN BREEDING SWINE

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

Each mL of Flunazine[®]-S (flunixin meglumine) Injectable Solution contains flunixin meglumine equivalent to 50 mg flunixin, 0.1 mg edetate disodium, 2.2 mg sodium formaldehyde sulfoxylate, 4.0 mg diethanolamine, 207.2 mg propylene glycol, 5.0 mg phenol as preservative, hydrochloric acid, water for injection q.s.

CLINICAL PHARMACOLOGY:

Flunixin meglumine is a potent non-narcotic, non-steroidal, analgesic agent with anti-inflammatory and antipyretic activity. It is significantly more potent than pentazocine, meperidine, and codeine as an analgesic in the rat yeast paw test.

Flunixin is known to persist in inflammatory tissues¹ and is associated with anti-inflammatory properties which extend well beyond the period associated with detectable plasma drug concentrations². Therefore, prediction of drug concentrations based upon estimated plasma terminal elimination half-life will likely underestimate both the duration of drug action and the concentration of drug remaining at the site of activity.

The pharmacokinetic profiles were found to follow a 2-compartmental model, although a deep (third) compartment was observed in some animals. The mean terminal elimination half-life (β half-life) of flunixin after a single intramuscular injection of flunixin (2.2 mg/kg) to pigs was between 3 and 4 hours. The mean observed maximum plasma concentration was 2944 ng/mL, achieved at a mean time of approximately 0.4 hours. The mean $AUC_{(0-100)}$ was 6431 ng*hr/mL. Following IM administration of flunixin, quantifiable drug concentration could be measured up to 18 hours post dose. The mean volume of distribution was 2003 mL/kg and the mean total clearance was 390 mL/hr/kg. The mean absolute bioavailability of flunixin following an intramuscular injection in the neck was 87%.

INDICATION:

Flunazine[®]-S (flunixin meglumine) Injectable Solution is indicated for the control of pyrexia associated with swine respiratory disease.

DOSE AND ADMINISTRATION:

The recommended dose for swine is 2.2 mg/kg (1 mg/lb; 2 mL per 100 lbs) body weight given by a single intramuscular administration. The injection should be given only in the neck musculature with a maximum of 10 mL per site.

Note: Intramuscular injection may cause local tissue irritation and damage. In an injection-site irritation study, the tissue damage did not resolve in all animals by Day 28 post-injection. This may result in trim loss of edible tissue at slaughter.

CONTRAINDICATIONS:

There are no known contraindications to this drug in swine when used as directed. Do not use in animals showing hypersensitivity to flunixin meglumine. Use judiciously when renal impairment or gastric ulceration is suspected.

RESIDUE WARNINGS:

Swine must not be slaughtered for human consumption within 12 days of the last treatment.

PRECAUTIONS:

As a class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, renal and hepatic toxicity. Sensitivity to drug-associated adverse events varies with the individual patient. Patients at greatest risk for adverse events are those that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be carefully approached. NSAIDs may inhibit the prostaglandins that maintain normal homeostatic function. Such prostaglandin effects may result in clinically significant disease in patients with underlying or pre-existing disease that has not been previously diagnosed.

Since many NSAIDs possess the potential to produce gastrointestinal ulceration, concomitant use of flunixin meglumine with other anti-inflammatory drugs, such as other NSAIDs and corticosteroids, should be avoided.

Not for use in breeding swine. The reproductive effects of Flunazine[®]-S (flunixin meglumine) Injectable Solution have not been investigated in this class of swine. Intramuscular injection may cause local tissue irritation and damage. In an injection-site irritation study, the tissue damage did not resolve in all animals by Day 28 post-injection. This may result in trim loss of edible tissue at slaughter.

SAFETY:

Flunixin was mildly irritating at the injection sites. No other flunixin-related changes (adverse reactions) were noted in swine administered a 1X (2.2 mg/kg; 1.0 mg/lb) dose for 9 days. Minimal toxicity manifested itself as statistically significant increased spleen weight at elevated doses (5X or higher daily for 9 days) with no change in normal microscopic architecture.

HOW SUPPLIED:

Flunazine[®]-S (flunixin meglumine) Injectable Solution, 50 mg/mL, is available in 100 mL multi-dose vial.

STORAGE:

STORE BETWEEN 2°C and 30°C (36°F and 86°F). KEEP FROM FREEZING.

REFERENCES

1. Lees P, Higgins AJ. Flunixin inhibits prostaglandin E₂ production in equine inflammation. *Res Vet Sci.* 1984; 37:347-349.
2. Odensvik K. Pharmacokinetics of flunixin and its effect on prostaglandin F_{2α} metabolite concentrations after oral and intravenous administration in heifers. *J Vet Pharmacol Ther.* 1995; 18:254-259.

To obtain an MSDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.

