

LincoMed® 100 & 300

Injection for Swine Antibiotic

(lincomycin, USP)

ANADA 200-368, Approved by FDA

DESCRIPTION

LincoMed® contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis* var. *lincolnensis*, which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

INDICATIONS

LincoMed® is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp.

It is also indicated for the treatment of mycoplasma pneumonia.

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1LIN019	100 mL	12
1LIN020	100 mL	12



BENEFITS

- **Accurate dosing:** Two concentrations for precise administration
- **Effective:** Contains the proven antibiotic lincomycin in injectable form
- **Versatile:** Use in all ages, at all stages of production
- **Economical:** Reduces losses due to profit-robbing arthritic damage
- **Safe:** Approved by FDA

RESIDUE WARNING

Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

See reverse side for Administration and Dosage.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com



LincoMed® 100 & 300

(lincomycin, USP)

Injection for Swine Antibiotic

ANADA 200-368, Approved by FDA

**FOR INTRAMUSCULAR USE IN SWINE.
FOR USE IN ANIMALS ONLY. NOT FOR HUMAN USE.
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.**

Restricted Drug (California) – Use Only as Directed

INDICATIONS:

LincoMed® is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp.

It is also indicated for the treatment of mycoplasma pneumonia.

CONTRAINDICATIONS:

As with all drugs, the use of LincoMed® is contraindicated in animals previously found to be hypersensitive to the drug.

RESIDUE WARNING:

Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

HUMAN WARNINGS:

Not for human use. Keep out of reach of children.

CAUTION:

If no improvement is noted within 48 hours, consult a veterinarian.

ADVERSE REACTIONS:

The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert to the possibility that it may occur. Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION:

For arthritis or mycoplasma pneumonia—5mg/lb body weight intramuscularly once daily for three to seven days as needed. One mL LincoMed® 100 per 20 lb body weight, or one mL LincoMed® 300 per 60 lb body weight, will provide 5 mg/lb.

For optimal results, initiate treatment as soon as possible. As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

EACH mL CONTAINS:

LincoMed® 100

Lincomycin hydrochloride equivalent to lincomycin 100 mg; also benzyl alcohol 9.45 mg added as a preservative.

LincoMed® 300

Lincomycin hydrochloride equivalent to lincomycin 300 mg; also benzyl alcohol 9.45 mg added as a preservative.

STORAGE:

**Store at controlled room temperature 20°C - 25°C (68°F - 77°F)
PROTECT FROM FREEZING.**

HOW SUPPLIED:

LincoMed® 100 and LincoMed® 300 (lincomycin injection, USP) are supplied in 100 mL (3.3 fl oz) multiple dose vials; 12 vials per case.

To obtain an MSDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.

