

Tetroxy[®] HCA-1772

(oxytetracycline HCl) Soluble Powder

FOR CONTROL AND TREATMENT OF
SPECIFIC DISEASES IN SWINE

Antibiotic

ANADA 200-144, Approved by FDA



MADE IN USA

INDICATIONS

For use in the drinking water of swine.

For the Control and Treatment of the Following Diseases in Swine: **Bacterial enteritis** caused by *Escherichia coli* and *Salmonella choleraesuis*, susceptible to oxytetracycline. **Bacterial pneumonia** caused by *Pasteurella multocida*, susceptible to oxytetracycline. **For Breeding Swine: Leptospirosis** (reducing the incidence of abortions and shedding leptospira) caused by *Leptospira pomona*, susceptible to oxytetracycline.

BENEFITS

- **Broad spectrum:** Economical treatment for a variety of common swine diseases
- **Convenient:** Zero withdrawal time before slaughter
- **Soluble:** Readily dissolves in water
- **Safe:** Approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TET015	1772 g (3.91 lb) pouch	8

CAUTION

Use as sole source of oxytetracycline. Not to be used for more than five consecutive days.

RESIDUE WARNING

Discontinue treatment 0 days prior to slaughter.

CONSULT MATERIAL SAFETY DATA SHEET
PRIOR TO USE.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@BimedaUS.com

See reverse side for Administration and Dosage.



Tetroxy® HCA-1772

(oxytetracycline HCl)
Soluble Powder

FOR CONTROL AND TREATMENT OF SPECIFIC DISEASES IN SWINE
Antibiotic

ANADA 200-144, Approved by FDA

NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

FOR USE IN DRINKING WATER ONLY
NOT FOR USE IN LIQUID TYPE B MEDICATED FEEDS

Restricted Drug (California) – Use only as directed.

ACTIVE INGREDIENT:

Each packet contains 648 g oxytetracycline HCl.
3.91 lbs. (1.77 kg) will treat 64,800 pounds of swine.

INDICATIONS:

For the Control and Treatment of the Following Diseases in Swine: Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*, susceptible to oxytetracycline. Bacterial pneumonia caused by *Pasteurella multocida*, susceptible to oxytetracycline.

For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding leptospira) caused by *Leptospira pomona*, susceptible to oxytetracycline.

DIRECTIONS FOR USE:

Mix fresh solutions daily. Use as sole source of drinking water. Do not mix product directly with milk or milk replacers. Administer one hour before or two hours after feeding milk or milk replacers. Administer up to five days to swine.

Treat swine for the indicated diseases at the rate of 10 mg/lb. body weight.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

CAUTION:

Use as sole source of oxytetracycline. Not to be used for more than five consecutive days in swine.

RESIDUE WARNING:

Discontinue treatment 0 days prior to slaughter. Consult material safety data sheet prior to use.

HOW SUPPLIED:

1772 g (3.91 lb) (1.77 kg) pouch; 8 pouches per case.

STORAGE:

Store below 77° F (25° C).

CONSULT MATERIAL SAFETY DATA SHEET PRIOR TO USE.

To obtain an MSDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.

