

Tetroxy[®] 343 (oxytetracycline HCl) Soluble Powder

A Broad Spectrum Antibiotic
343 g oxytetracycline HCl per pound

ANADA 200-247, Approved by FDA



MADE IN USA

INDICATIONS

For the control and treatment of specific diseases in poultry, cattle, sheep, swine and bees. For the marking of skeletal tissues in finfish fry and fingerlings as an aid in identification.

BENEFITS

- **Versatile:** Approved for use in multiple food animal species – including honey bees and fish
- **Broad spectrum:** Controls a variety of disease organisms
- **Concentrated:** Each pound of product contains 343 g oxytetracycline HCl
- **Economical:** Each 135.5 g incremental pack treats 10,240 lb of swine, cattle or sheep at 10 mg/lb of body weight
- **Short withdrawal time:** Only five days after the last treatment in turkeys, cattle and sheep, zero days in swine
- **Convenient:** Four package sizes available to meet the varying demands of producers
- **Oral dosage form:** Reduces handling and animal stress
- **Safe:** Approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TET040	135.5 g (4.78) pouch	25
1TET041	272.2 g (9.6 oz) pouch	25
1TET042	907.2 g (2 lb) pail	1
1TET043	2.26 kg (5 lb) pail	1

RESIDUE WARNINGS

Use as sole source of oxytetracycline. Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. An additional withdrawal time beyond the grow-out period is not needed for fish. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com

See reverse side for Administration and Dosage.



BimedaUS.com

Tetroxy® 343 (oxytetracycline HCl) Soluble Powder

A Broad Spectrum Antibiotic • 343 g oxytetracycline HCl per pound

ANADA 200-247, Approved by FDA

For oral use only in poultry, cattle, sheep, swine and bees. Fish should be treated by immersion.

FOR USE IN DRINKING WATER ONLY.

NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

FOR ANIMAL USE ONLY. KEEP OUT OF REACH OF CHILDREN.

Restricted Drug(s) (California). Not For Human Use. Use Only As Directed.

INDICATIONS AND DIRECTIONS FOR USE:

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon. The enclosed measuring cup for the 2 lb and 5 lb sizes holds 68 g (2.4 oz) of product when leveled at the 80 mL mark.

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage. See Residue Warnings.

ACTIVE INGREDIENT:

135.5 g (4.78 oz) Pouch

This packet contains 102.4 grams of oxytetracycline HCl.

This packet will make:

- 512 gallons (1,938 L) containing 200 mg oxytetracycline HCl per gallon
- 256 gallons (969 L) containing 400 mg oxytetracycline HCl per gallon
- 128 gallons (484.5 L) containing 800 mg oxytetracycline HCl per gallon

272.2 g (9.6 oz)

This packet contains 204.8 grams of oxytetracycline HCl.

This packet will make:

- 1,024 gallons (3,876 L) containing 200 mg oxytetracycline HCl per gallon (equivalent to 52.8 mg/L)
 - 512 gallons (1,938 L) containing 400 mg oxytetracycline HCl per gallon (equivalent to 105.7 mg/L)
 - 256 gallons (969 L) containing 800 mg oxytetracycline HCl per gallon (equivalent to 211.4 mg/L)
- This packet will treat 20,480 pounds of swine, cattle or sheep at 10 mg/pound.

907.2 g (2 lb) Pail

This pail contains 686 grams of oxytetracycline HCl and will treat 68,600 pounds of swine, cattle or sheep at 10 mg/pound

2.26 kg (5 lb) Pail

This pail contains 1,715 grams of oxytetracycline HCl and will treat 171,500 pounds of swine, cattle or sheep at 10 mg/pound

DISEASE		DOSAGE	135.5 g (4.78 oz) PACKETS/ 2 GALLONS STOCK SOLUTION	272.2 g (9.6 oz) PACKETS/ 2 GALLONS STOCK SOLUTION	907.2 g (2 lb) & 2.26 kg (5 lb) SCOOPS/2 GALLONS STOCK SOLUTION
CHICKENS	Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	200-400 mg/gal	½ - 1	¼ - ½	1 - 2
	Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> .	400-800 mg/gal	1 - 2	½ - 1	2 - 4
	Fowl cholera caused by <i>Pasteurella multocida</i> .	400-800 mg/gal	1 - 2	½ - 1	2 - 4
TURKEYS	Hexamitiasis caused by <i>Hexamita meleagridis</i> .	200-400 mg/gal	½ - 1	¼ - ½	1 - 2
	Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	400 mg/gal	1	½	2
	Growing Turkeys—Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis).	25 mg/lb body weight daily	Varies with age & water consumption (1 packet will treat 4,096 pounds of turkeys.)	Varies with age & water consumption (1 packet will treat 8,192 pounds of turkeys.)	Varies with age & water consumption (1 pound will treat 13,720 pounds of turkeys.)

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

DISEASE	DOSAGE
SWINE	
Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia caused by <i>Pasteurella multocida</i> .	
For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> .	
CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE	
Bacterial enteritis caused by <i>Escherichia coli</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	
SHEEP	
Bacterial enteritis caused by <i>Escherichia coli</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	
HONEY BEES	
For control of American Foulbrood caused by <i>Paenibacillus larvae</i> , and European Foulbrood caused by <i>Streptococcus pluton</i> susceptible to oxytetracycline.	200 mg/colony Administer in 3 applications of sugar syrup or 3 dustings at 4 to 5 day intervals. Apply dust on the outer parts or ends of the frames. See Residue Warnings.
CONTRAINDICATION: Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.	
FISH	
For the marking of skeletal tissues in finfish fry and fingerlings as an aid in identification.	Immerse fry or fingerling finfish in 200 to 700 mg OTC (buffered)/liter of water for 2 - 6 hours. Solution should be tested on a small number of fish before full-scale use. Once mixed with water, solution should be used immediately. See Residue Warnings.

ADVERSE REACTIONS: Oxytetracycline HCl will acidify the water. The pH should be maintained at an acceptable level for fish by the addition of a buffer. Monitor water quality and temperature.

DISPOSAL OF TREATED WATER: Do not discharge marking immersion water containing oxytetracycline into surface waters.

PRECAUTION: Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

RESIDUE WARNINGS:

Use as sole source of oxytetracycline. Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. An additional withdrawal time beyond the grow-out period is not needed for fish. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Manufactured by:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

2296-15 Tetroxy 343 TS 04/15

TAKE TIME OBSERVE LABEL DIRECTIONS

BimedaUS.com