

Tetroxy[®] 25 (oxytetracycline HCl) Soluble Powder

A Broad Spectrum Antibiotic
25 g oxytetracycline HCl per pound

ANADA 200-146, Approved by FDA



MADE IN USA

INDICATIONS

For control and treatment of specific diseases (see reverse side) in poultry, cattle, sheep, swine, and honey bees.

BENEFITS

- **Versatile:** Approved for use in multiple food animal species
- **Broad spectrum:** Treats and controls a variety of disease organisms
- **Convenient:** Three package sizes available to meet the varying demands of producers
- **Short withdrawal time:** Only five days after the last administration
- **Oral dosage form:** Reduces handling and animal stress
- **Soluble:** Readily dissolves in water
- **Easy measuring:** Bulk sizes include handy measuring scoop
- **Safe:** Approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TET044	181.5 g (6.4 oz) pouch	25
1TET045	907.2 g (2 lb) pail	1
1TET046	2.26 kg (5 lb) pail	1

PRECAUTION

Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

RESIDUE WARNINGS

Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.



See reverse side for Administration and Dosage.

Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com

BimedaUS.com

Tetroxy® 25 (oxytetracycline HCl) Soluble Powder

A Broad Spectrum Antibiotic • 25 g oxytetracycline HCl per pound

ANADA 200-146, Approved by FDA

For control and treatment of specific diseases in poultry, cattle, sheep and swine and honey bees.

FOR USE IN DRINKING WATER ONLY.

NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN.

Restricted Drug (California) - Use Only as Directed

INDICATIONS AND DIRECTIONS FOR USE:

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

ACTIVE INGREDIENT:

181.5 g (6.4 oz) Pouch

Contains 10 grams of oxytetracycline HCl and will treat 1,000 pounds of swine, cattle or sheep at 10 mg/pound

907.2 g (2 lb) Pail

Contains 50 grams of oxytetracycline HCl and will treat 5,000 pounds of swine, cattle or sheep at 10 mg/pound

2.26 kg (5 lb) Pail

Contains 125 grams of oxytetracycline HCl and will treat 12,800 pounds of swine, cattle or sheep at 10 mg/pound

DISEASE		DOSAGE	181.5 g (6.4 oz) PACKETS PACKETS/2 GALLONS STOCK SOLUTION	907.2 g (2 lb) PAIL POUNDS/2 GALLONS STOCK SOLUTION	2.26 kg (5 lb) PAIL POUNDS/2 GALLONS STOCK SOLUTION
CHICKENS	Infectious synovitis caused by <i>Mycoplasma synoviae</i>	200-400 mg/gal	5-10	2 - 4	2 - 4
	Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i>	400-800 mg/gal	10-20	4 - 8	4 - 8
	Fowl cholera caused by <i>Pasteurella multocida</i>	400-800 mg/gal	10-20	4 - 8	4 - 8
TURKEYS	Hexamitiasis caused by <i>Hexamita meleagridis</i>	200-400 mg/gal	5-10	2 - 4	2 - 4
	Infectious synovitis caused by <i>Mycoplasma synoviae</i>	400 mg/gal	10	4	4
	Growing Turkeys—Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis)	25 mg/lb body weight daily	Varies with age & water consumption (1 packet will treat 400 pounds of turkeys.)	Varies with age & water consumption (1 pound will treat 1,000 pounds of turkeys.)	Varies with age & water consumption (1 pound will treat 1,000 pounds of turkeys.)

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage. See Residue Warnings.

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

DISEASE	DOSAGE
SWINE	
Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	
For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> .	
CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE	
Bacterial enteritis caused by <i>Escherichia coli</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	
SHEEP	
Bacterial enteritis caused by <i>Escherichia coli</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	
HONEY BEES	
For control of American Foulbrood caused by <i>Paenibacillus larvae</i> , and European Foulbrood caused by <i>Streptococcus pluton</i> susceptible to oxytetracycline.	200 mg/colony Administer in 3 applications of sugar syrup or 3 dustings at 4 to 5 day intervals. Apply dust on the outer parts or ends of the frames. See Residue Warnings.
CONTRAINDICATION: Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.	

PRECAUTION:

Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

RESIDUE WARNINGS:

Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Manufactured by:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

