

EXODUS[®]

(pyrantel pamoate) Paste

DATA SHEET



INDICATIONS

For the removal and control of mature infections in horses and ponies:

large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*)

small strongyles

pinworms (*Oxyuris equi*)

large roundworms (*Parascaris equorum*)

BENEFITS

- ✓ **Apple flavored**
- ✓ **Proven:** Contains pyrantel pamoate for safe worm removal and control
- ✓ **Versatile:** Over 90% effective against most major equine worms
- ✓ **Convenient:** Pre-measured plastic syringes provide accurate dosing
- ✓ **Easy-to-Use:** Available in two size syringes



LIST No	UNIT PACKAGE	CASE SIZE
1EXO001	23.6 g	72 (6x12)
1EXO010	23.6 g	100
1EXO002	47.2 g	36

See reverse for Administration & Dosage

Exodus[®] (pyrantel pamoate) Paste




DESCRIPTION:

Exodus (pyrantel pamoate) Paste is a pale yellow to buff paste containing 43.9% w/w pyrantel pamoate in an inert vehicle. Each single dose syringe contains 3.6 grams of pyrantel base in 23.6 grams paste. Each multi-dose syringe contains 7.2 grams of pyrantel base in 47.2 grams paste. Each mL contains 171 mg pyrantel base as pyrantel pamoate.

COMPOSITION:

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below.



Chemical name:
(E)-1,4,5,6-tetrahydro-1-methyl-2-[2-(2-thienyl)-vinyl]-pyrimidine 4,4'-methylenebis [3-hydroxy-2-naphtholate] (1:1)

INDICATIONS FOR USE:

For the removal and control of mature infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles; pinworms (*Oxyuris equi*); and large roundworms (*Parascaris equorum*) in horses and ponies.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND TREATMENT:

Exodus (pyrantel pamoate) Paste is to be administered as a single oral dose of 3 mg pyrantel base per lb of body weight. The single dose syringe has 4 weight mark increments. The multi-dose syringe has 8 weight mark increments. Each weight mark indicates the recommended dose for 300 lb of body weight. One-half of the multi-dose syringe treats a 1200 lb horse.

EXODUS PASTE DOSAGE		
Body Weight Range	Volume	mg Pyrantel Base
up to 300 lb	1/4 syringe	900 mg
301 - 600 lb	1/2 syringe	1800 mg
601 - 900 lb	3/4 syringe	2700 mg
901 - 1200 lb	1 full syringe	3600 mg

EXODUS MULTI-DOSE DOSAGE		
Body Weight Range	Volume	mg Pyrantel Base
up to 300 lb	1/8 syringe	900 mg
301 - 600 lb	1/4 syringe	1800 mg
601 - 900 lb	3/8 syringe	2700 mg
901 - 1200 lb	1/2 syringe	3600 mg

Note: Position ring-gauge over appropriate mark on plunger. Each mL contains 171 mg of pyrantel base as pyrantel pamoate.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

For maximum control of parasitism, it is recommended that foals (2-8 months of age) be dosed every 4 weeks. To minimize the potential source of infection that the mare may pose to the foal, the mare should be treated 1 month prior to anticipated foaling date followed by retreatment 10 days to 2 weeks after birth of foal. Horses and ponies over 8 months of age should be routinely dosed every 6 weeks.

ADMINISTRATION:

After removing the cap, the paste should be deposited on the dorsum of the tongue. Introduce the nozzle end of the syringe at the corner of the mouth. Direct the syringe backwards and depress the plunger to deposit the paste onto the tongue. Given in this manner, it is unlikely that rejection of the paste will occur. Raising the horse's head sometimes assists in the swallowing process. When only part of the paste has been used, replace the cap on the syringe nozzle.

EFFICACY:

Critical (worm count) studies in horses demonstrated that pyrantel pamoate paste administered at the recommended dosage was efficacious against mature infections of *Strongylus vulgaris* (>90%), *S. edentatus* (69%), *S. equinus* (>90%), *Oxyuris equi* (81%), *Parascaris equorum* (>90%), and small strongyles (>90%).

WARNINGS

For oral animal use only. Do not use in horses intended for human consumption.

HUMAN WARNINGS:

Keep out of the reach of children.

It is recommended that severely debilitated animals not be treated with this preparation.

OTHER WARNINGS:

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

RECOMMENDED STORAGE:

Store at 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 30°C (59°F - 86°F).

DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS

Any unused product or waste materials should be disposed of in accordance with local requirements.

Approved by FDA under ANADA # 200-350

MANUFACTURED FOR:

Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

N.A. CORP. ADDRESS:

Bimeda, Inc.
One Tower Lane, Oakbrook Terrace, IL 60181

For further information and queries, please contact Bimeda at (888) 524-6332 or US-Info@Bimeda.com.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

www.BimedaUS.com