

LevaMed™ (levamisole hydrochloride) Soluble Pig Wormer



FOR USE IN DRINKING WATER

Anthelmintic

RECOMMENDATIONS

LevaMed (levamisole hydrochloride) Soluble Pig Wormer is a broad-spectrum anthelmintic and is effective against the following nematode infections in swine.

Large Roundworms: (*Ascaris suum*)

Nodular Worms: (*Oesophagostomum* spp.)

Lungworms: (*Metastrongylus* spp.)

Intestinal Threadworms: (*Strongyloides ransomi*)

DIRECTIONS FOR PREPARING SOLUBLE PIG WORMER SOLUTION

When you are ready to worm your pigs, add water to the powder in this bottle up to the 500 ml mark. Agitate to mix thoroughly before using. If any solution is left over, it may be stored at up to 25°C for up to 3 months in this tightly capped bottle; agitate well before using.

DIRECTIONS FOR USE

Withholding water from pigs prior to treatment is not necessary for optimum anthelmintic efficacy and is not recommended during hot weather. Add 10 ml (2 teaspoonfuls) of the solution from this bottle to 1 gallon of water; mix thoroughly. Allow one gallon of medicated water for each 100 pounds body weight of pigs to be treated. No other source of water should be offered. As soon as pigs have consumed all the medicated water, resume use of regular water.

This bottle contains 18.15 grams of levamisole hydrochloride activity which will treat the following:

200 - 25 lb pigs

100 - 50 lb pigs

50 - 100 lb pigs

25 - 200 lb pigs

NOTE: Careful estimates of pig weights are essential for proper performance of this product.

Pigs maintained under conditions of constant worm exposure may require retreatment within 4-5 weeks after the first treatment due to reinfection.

OTHER WARNINGS

Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

CAUTION

Consult your veterinarian before administering LevaMed to sick swine. **Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.**

Salivation or muzzle foam may be observed. This reaction is occasionally seen and will disappear in a short time after medication.

If pigs are infected with mature lungworms, coughing and vomiting may be observed soon after medicated water is consumed. This reaction is due to the expulsion of worms from the lungs and will be over in several hours. Follow recommended dosage carefully to assure removal of worms and avoid an overdose of LevaMed solution.

RESIDUE WARNING

Do not administer within 72 hours of slaughter for food.

STORAGE

Store at 20°C - 25°C (68°F to 77°F), excursions permitted between 15°C - 30°C (59°F and 86°F).

NOT FOR USE IN HUMANS.

KEEP OUT OF REACH OF CHILDREN.

Restricted Drug (California) - Use Only as Directed

Approved by FDA under ANADA # 200-313

List Number	Pack Size	Case Size
1LEV003	20.17 g	12

MANUFACTURED BY:

Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

N.A. CORP. ADDRESS:

Bimeda, Inc.
One Tower Lane
Oakbrook Terrace, IL 61081

For further information and queries, contact Bimeda at (888) 524-6332 or US-info@Bimeda.com