

PenAqua Sol-G[®]

(penicillin G potassium)
Antibiotic for drinking water



For Oral Use in Turkeys Only
For Veterinary Use Only

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

INDICATIONS

For the treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

ACTIVE INGREDIENT:

313 gram (11 ounce) Pouch – 0.500 Billion Units penicillin G potassium – Nonsterile
626 gram (22 ounce) Pouch – 1 Billion Units penicillin G potassium – Nonsterile

DOSAGE AND ADMINISTRATION

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS

0.500 billion I.U. Pouch

Combine contents and approximately 1-1/2 pints (710 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems – Pour the concentrated solution into a glass or plastic container then add enough water to make 2.6 gallons (9.9 liters) of stock solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water]. The automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

Gravity Flow Watering Systems – Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

1.000 billion I.U. Pouch

Combine contents and approximately 3 pints (1420 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems – Pour the concentrated solution into a glass or plastic container then add enough water to make 5.2 gallons (19.7 liters) of stock solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water]. The automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

PenAqua Sol-G is a registered trademark of Bimeda Inc.
Rev. 8/24 M-24-0570

Gravity Flow Watering Systems – Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1.000 billion units of penicillin G per gallon (3.8 liters).

WARNINGS

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.
Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.

STORAGE

Store at room temperature between 20° - 25°C (68° to 77°F), with excursions up to 40°C (104°F).

| PRODUCT NO. | UNIT SIZE | CASE SIZE |
|-------------|-----------------------------------|-----------|
| 1PEN405 | 0.5 B.U. pouch (Net Wt: 313 g) | 20 |
| 1PEN406 | 1.0 B.U. pouch (Net Wt: 626 g) | 8 |

Approved by FDA under ANADA # 200-372

Keep Out of Reach of Children
Restricted Drug (California) – Use Only as Directed

MANUFACTURED FOR:

Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

N.A. CORP. ADDRESS:

Bimeda, Inc.
475 N. Martingale, Suite 1200
Schaumburg, IL 60173