

# TETROXY<sup>®</sup> HCA-1400 (oxytetracycline hydrochloride)

# DATA SHEET



## INDICATIONS

Tetroxy<sup>®</sup> HCA-1400 Soluble Powder is indicated for the control and treatment of:

**Chicken** - infectious synovitis caused by *Mycoplasma synoviae*. Chronic respiratory disease (CRD) and air sac infections caused by *Mycoplasma gallisepticum* and *Escherichia coli*. Fowl cholera caused by *Pasteurella multocida*.

**Turkeys** - hexamitiasis caused by *Hexamita meleagridis*. Infectious synovitis caused by *Mycoplasma synoviae*. Growing Turkeys - Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis).

**Swine** - bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*. Bacterial pneumonia caused by *Pasteurella multocida*.  
**For Breeding Swine:** Leptospirosis caused by *Leptospira pomona*.

## BENEFITS

- ✓ **Versatile:** Antibiotic treatment for chickens, turkeys and swine
- ✓ **Zero withdrawal time** before slaughter
- ✓ **Soluble:** Readily dissolves in water
- ✓ **Approved by FDA**

LIST No	UNIT PACKAGE	CASE SIZE
1TET014	1400 g	10



**Caution:** Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

See reverse for Administration & Dosage

# Tetroxy® HCA-1400

## (oxytetracycline hydrochloride)

### Soluble Powder



Approved by FDA under ANADA # 200-144

**Antibiotic**

For control and treatment of specific diseases in poultry and swine.

**FOR USE IN DRINKING WATER ONLY.**

**NOT FOR USE IN LIQUID MEDICATED FEEDS.**

**NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN.**

**Restricted Drug (California) - Use Only as Directed**

**CAUTION:**

Federal (U.S.A) law restricts this drug to use by or on the order of a licensed veterinarian.

**INDICATIONS AND DIRECTIONS FOR USE:**

Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	DISEASE	TREATMENT LEVEL	GALLONS STOCK/PACKET
<b>C H I C K E N S</b>	Infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	20 - 10
	Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
	Fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
<b>T U R K E Y S</b>	Hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	20 - 10
	Infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	<b>GROWING TURKEYS</b> - Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
<b>S W I N E</b>	<b>FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE:</b> Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	<b>FOR BREEDING SWINE:</b> Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

**DIRECTIONS FOR USE:**

Mix fresh solutions daily. Use as sole source of drinking water-do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. Consult a poultry diagnostic laboratory or poultry pathologist for diagnosis and advice on dosage for chickens and turkeys. As a generalization, 200 chickens will drink one gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer 7 to 14 days to chickens & turkeys and up to 5 days to swine.

**NOTE:**

The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

**PRECAUTION:**

Use as sole source of oxytetracycline. Not to be used for more than 14 consecutive days in chickens and turkeys or 5 consecutive days in swine.

▶ **RESIDUE WARNING:** Do not feed to birds producing eggs for human consumption. Discontinue treatment of chickens, turkeys and swine 0 days prior to slaughter ◀

**RECOMMENDED PACKET STORAGE CONDITIONS:**

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

**CONSULT SAFETY DATA SHEET PRIOR TO USE.**

**DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS**

Any unused product or waste materials should be disposed of in accordance with local requirements.

**MANUFACTURED BY:**

Bimeda, Inc.  
Le Sueur, MN 56058

**N.A. CORP. ADDRESS:**

Bimeda, Inc.  
One Tower Lane, Oakbrook Terrace, IL 60181

For further information and queries, please contact Bimeda at (888) 524-6332 or US-Info@Bimeda.com.

TAKE TIME



OBSERVE LABEL DIRECTIONS

[www.BimedaUS.com](http://www.BimedaUS.com)

Tetroxy® is a registered trademark of Bimeda, Inc.  
Tetroxy® HCA-1400 TS 11/19

Global Excellence in Animal Health

