

# TETROXY<sup>®</sup> HCA-1772

(oxytetracycline hydrochloride)

# DATA SHEET



## INDICATIONS

Tetroxy<sup>®</sup> HCA-1772 Soluble Powder is indicated for the control and treatment of the following disease in swine:


Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*.

Bacterial pneumonia caused by *Pasteurella multocida*.

**For Breeding Swine:** Leptospirosis caused by *Leptospira pomona*.

## BENEFITS

- ✓ Zero withdrawal time before slaughter
- ✓ Soluble: Readily dissolves in water
- ✓ Approved by FDA



LIST No	UNIT PACKAGE	CASE SIZE
1TET015	1772 g	8




**Caution:** Federal (U.S.A) law restricts this drug to use by or on the order of a licensed veterinarian.

**See reverse for Administration & Dosage**

# Tetroxy<sup>®</sup> HCA-1772

(oxytetracycline hydrochloride)

## Soluble Powder



Approved by FDA under ANADA # 200-144

**Antibiotic**

For control and treatment of specific diseases in swine.

**FOR USE IN DRINKING WATER ONLY.**

**NOT FOR USE IN LIQUID MEDICATED FEEDS.**

**NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN.**

**Restricted Drug (California) - Use Only as Directed**

**CAUTION:**

Federal (U.S.A) law restricts this drug to use by or on the order of a licensed veterinarian.

**INDICATIONS:**

	DISEASE	TREATMENT LEVEL
SWINE	<b>FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE:</b> Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline. <b>FOR BREEDING SWINE:</b> Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight

**DIRECTIONS FOR USE:**

Mix fresh solutions daily. Use as sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. Administer up to 5 days to swine.

**NOTE:**

The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

**PRECAUTION:**

Use as sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine.

► **RESIDUE WARNING:** Discontinue treatment of swine 0 days prior to slaughter ◀

**RECOMMENDED PACKET STORAGE CONDITIONS:**

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

**CONSULT SAFETY DATA SHEET PRIOR TO USE.**

**DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS**

Any unused product or waste materials should be disposed of in accordance with local requirements.

**MANUFACTURED BY:**

Bimeda, Inc.  
Le Sueur, MN 56058

**N.A. CORP. ADDRESS:**

Bimeda, Inc.  
One Tower Lane, Oakbrook Terrace, IL 60181

For further information and queries, please contact Bimeda at (888) 524-6332 or US-Info@Bimeda.com.

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

[www.BimedaUS.com](http://www.BimedaUS.com)

Tetroxy<sup>®</sup> is a registered trademark of Bimeda, Inc.  
Tetroxy<sup>®</sup> HCA-1772 TS 11/19

Global Excellence in Animal Health

 **Bimeda<sup>®</sup>**