

0.5 mg/mL Glucocorticoid

Bimasone is recommended for the various rheumatic, allergic, dermatologic, and other disease states which are known to be responsive to the anti-inflammatory corticoids.

Flumethasone has the highest relative glucocorticoid activity (120) and long duration of effect compared to other commonly used corticosteroids.

*FluCort® is a registered trademark of Zoetis, Inc. Federal law restricts this drug to use by or in the order of a licensed veterinarian. M-24-0493



Learn more at www.bimedaus.com/bimasone



RELATIVE POTENCIES OF COMMONLY USED CORTICOSTEROIDS

COMPOUND	RELATIVE GLUCOCORTICOID ACTIVITY	RELATIVE MINERALOCORTICOID ACTIVITY	DURATION OF EFFECT ^A (ALCOHOL FORM)
Flumethasone	120	0	L
Betamethasone	25	0	L
Dexamethasone	25	0	L
Isoflupredone	25	25	L
Triamcinolone acetonide	30	0	I
Triamcinolone	5	0	I
Fludrocortisone	10	125	I
Methylprednisolone	5	0.5	1
Prednisolone	5	0.8	I
Prednisone	5	0.8	I
Cortisol	1	1	S

^{*}Reprinted from Merck Veterinary Manual.

have it for when you need it most.



